

A Study on Nosocomial Infections

Mahesh Sharma¹

Available online at: www.xournals.com

Received 6th September 2018 | Revised 19th October 2018 | Accepted 24th December 2018

Abstract:

In current scenario, Nosocomial infections are caused by different factors such as inappropriate hygiene. Patients in hospital can easily get infections of diseases like methicillin resistant staphylococcus aureus (MRSA) from hospital staff who do not practice basic hygienic measures. Nosocomial infection may also be result of contaminated use of injections, inappropriately bandaged incisions during surgeries, contaminated surgical equipment's, catheters, organ transplant and etc. However, frequency of these type of cases have become less in number today due to advanced technology but still it's the primary role of hospital staff to cure their patients from any suspected nosocomial infection. Here in this paper an effort has been done to understand the nosocomial infection and preventive measure taken to reduce the risk given by World Health Organization.

Keywords: Nosocomial Infections, World Health Organization, Preventive Measures

Authors:

1. Maharaja Ranjit Singh College of Professional Sciences, Indore, Madhya Pradesh, INDIA

Introduction

World Health Organization reported that around 109 people are hospitalized yearly for different reasons and 10 million people are infected with nosocomial infection. “Nosocomial” comes from Greek word *nosokomos*, means nosos=disease and komien= to care for (Mourud, 2010). Nosocomial Infection or Hospital Acquired Infection or Health Care-associated Infections can be defined as “An infection acquired in the hospital by a patient who was admitted for a reason other than that infection. An infection occurring in a patient in a hospital or other healthcare facility in whom the infection was not present or incubating at the time of admission. This includes infections acquired in the hospital but appearing after discharge, and also occupational infections among the staff of the facility” (<http://apps.who.int>). As per the survey, it was found that the patients in intensive care unit (ICU) are more prone to acquire the nosocomial infection in comparison to other ward patients (Özer, *et al.*, 2015; Patel, *et al.*, 2013).

Microorganisms such as a virus, bacteria, fungus etc. cause nosocomial infections. Different factors which are responsible for nosocomial infections are high density of pathogens and population in hospital, immunity of patients, poor hygiene, inadequate cleaning, increased number of medical procedures, etc. (Fang, 2012).

Robert A Weinstein, in his research paper, represents a complete research addressing to the changes in the medical science with respect to nosocomial infections in either of the way. Further, in his study, he also showed the noteworthy impact of progression in technology in medical and healthcare in relation to nosocomial infections. Jessica Lietz concluded in her research article by giving the general overview about the nosocomial infections focusing more on the general arguments. National Center for Infectious Diseases carried out a comprehensive research on the nosocomial infections in the United States discussing the key components of the infections, further comparing the rate of infections in the urban and rural setting. Toni Rizzo highlights on the common routes of infections in hospital i.e. respiratory procedures, intravenous (IV) procedures, surgery and wound and urinary bladder catheterization.

Types of Nosocomial Infections

Approximately 80% of all nosocomial infections occur at following four frequent sites:

a) Central Line-Associated Bloodstream Infections (CLABSI)

Prolonged used of Catheters in central line can cause serious bloodstream infections. According to the survey, these type of infections are the most deadly nosocomial infections with the death occurrence rate of 12%–25%. In United State hospitals, the decrease of 46% in CLABSI was seen from 2008 to 2013 but still, 30,100 CLABSI occur in ICU and acute facilities wards every year. In this, coagulase-negative staphylococci are the most common type of pathogens, accounting for 27.9%.

b) Catheter-Associated Urinary Tract Infections (CAUTI)

This type of infection is mainly caused by endogenous microflora and is the most frequently diagnosed, approximately 36% of the 2 million infections that occur yearly. In 2011, acute care hospitals stats shows that UTIs account for more than 12% of reported infections. It can further develop to serious complications in all patients, for instance, epididymitis and prostatitis in males, and pyelonephritis etc. *E. coli* (25.8%), enterococcus (15.9%), and *P. aeruginosa* (12%) are the most common type of nosocomial pathogens.

c) Surgical Site Infections (SSI)

This type of infections are the second most common type of nosocomial infections (approximately 17%) caused by *Staphylococcus aureus* which will ultimately result in death or hospitalization. It has been classified by level of contamination present at the time of surgery i.e. clean (class I), clean-contaminated (class II), contaminated (class III), and dirty (class IV) procedures since 1964. Pathogens which causes surgical site infections arise from endogenous microflora of the patient.

d) Ventilator-Associated Pneumonia (VAP)

Ventilator-associated pneumonia usually occur within 48 h after the tracheal incubation and found in 9–27% of patients on mechanically assisted

ventilator. Common symptoms of ventilator-associated pneumonia are fever, leucopenia, and bronchial sounds etc. *P. aeruginosa* (16.9%), *S. aureus* (16.1%), and *Enterobacter sp.* (10.5%) are most common pathogen causing nosocomial pneumonia. (Root, *et al.* 63; Khan, *et al.*, 2017; Mohammed, *et al.* 2014).




Figure 1: Sites of the most common Nosocomial Infections (Source: <http://apps.who.int>)

Nosocomial Pathogens

The infecting pathogens differ between different patient, different medical amenities, different health care settings, and different countries according to the World Health Organization 2002 guidelines. Nosocomial infections are mainly caused by common pathogens found in general. Following are the organisms which can cause nosocomial infections:

- a) Bacteria
- b) Viruses
- c) Fungi
- d) Parasites

In all pathogens bacteria are responsible for about 90% nosocomial infections. Bacteria are the most widely occurring nosocomial pathogens and their reservoir can be endogenous or exogenous, in which endogenous bacteria are from patient’s own flora whereas exogenous are from another patient, hospital staff. Further, difference in bacteria can be made among commensal bacteria (opportunistic) and pathogenic bacteria, in which commensal bacteria are part of normal flora of healthy women whereas pathogenic bacteria have greater virulence, and cause infections

irrespective of host status. Intravascular line infection and Urinary infection are caused by cutaneous coagulase-negative staphylococci and *Escherichia coli* respectively are the best example of commensal bacteria. Anaerobic Gram-positive rods (e.g. *Clostridium*) cause gangrene is example of pathogenic bacteria.

Other than bacteria, viruses are also common pathogen causing nosocomial infection and are generally transmitted by parenteral route i.e. transfusion, injections, dialysis or endoscopy or by faecal-oral route. Hepatitis A, B and C, RSV, Rotavirus, Enterovirus, CMV, HIV, etc. are different type of viruses causing nosocomial infections.

Fungi and parasites pathogens causes nosocomial infections during extended antibiotic therapy. In this fungal infections have become common cause of nosocomial infection. *Giardia lamblia* is an example of a parasite that can be easily spread among adults or children. Candidiasis is the most common type of nosocomial fungal infection.

25% of all these infections can be prevented, if hospital people start taking proper precautions. Below table 1 shows the site of infection and common pathogens causing nosocomial infection (<http://apps.who.int>; Khan, *et al.* 2017; Mourud, 2010).

Table 1: Site of Infection and Common Pathogens

Location	Pathogens
Blood Stream	<i>Coagulase-negative staphylococci</i> (CNS) <i>S. aureus</i> <i>P. aeruginosa</i> <i>Candida sp</i>
Pneumonia	CNS <i>S. aureus</i> <i>P. aeruginosa</i> Respiratory syncytial virus
Skin/Soft Tissue/Surgical Site	CNS <i>S. aureus</i>
Gastrointestinal Tract	Rotavirus
Conjunctivitis/Occular	CNS <i>P. aeruginosa</i>
Urinary Tract	Gram-negative bacilli

		<i>Enterococci</i>
Endocarditis		CNS <i>S. aureus</i>
Central Nervous System		CNS <i>S. aureus</i>
Osteoarthritis		<i>S. aureus</i> Group B Streptococci

(Source: Mohammed, *et al.* 2014)

Mode of Transmission

Nosocomial pathogens are transmitted in hospital by numerous route and many a time same pathogens may be transmitted by more than one route also. Contact, airborne, droplet route, common vehicle and vector borne are main route of transmission (Nazir and Kadri, 2014).

Prevention of Nosocomial Infections

Every individual in hospitals are individually responsible for prevention of nosocomial infections.

Management of hospitals should be done in such a way that hygiene is maintained always. There are different national or regional and hospitals programs to control the nosocomial infection.

National or Regional Programmes

These programs are developed by responsible authority to support the hospitals in reducing the risk and these programs must:

- Develop a national system to monitor infections.
- Develop and continually update guidelines.
- Provide facility to access materials and products required for hygiene and safety.
- Provide training programmes for health care professionals.
- Set a national objectives

Table 2: Infection Control Responsibility

Role	Responsibility
Role of Hospital Management	<ul style="list-style-type: none"> • Establish an Infection Control Committee. • Monitor infections and application of preventive measures • Education and training of all hospital staff. • Occasionally reviewing the status of nosocomial infections.
Role of the Physician	<ul style="list-style-type: none"> • Protect their own patients from other infections • Inform cases of nosocomial infection to the team and infected patients • Guide patients, visitors and hospital staff how to prevent from nosocomial infection.
Role of the Microbiologist	<ul style="list-style-type: none"> • Handle patient and staff specimens for microbiological diagnosis • Develop proper guidelines for collection, transport, and handling of microbiological specimens • Inform Infection Control Committee about the results.
Role of the Nursing Staff	<ul style="list-style-type: none"> • Take part in the Infection Control Committee • Develop training programmes for members of the nursing staff • Safe and adequate supply of ward equipment, drugs and patient care supplies. • Identify nosocomial infections • Self hygiene
Role of the Food Service	<ul style="list-style-type: none"> • To maintain the high level safety follow the proper criteria for the purchase of foodstuffs, equipment use, and cleaning procedures. • Clean all equipment used, working and storage areas. • Give proper instructions for handwashing, clothing, staff responsibilities and daily disinfection duties

(Source: <http://apps.who.int>)

Conclusion

After the detailed study about the nosocomial infection, it was found that the nosocomial infection is increasing day by day and caused just because of the inappropriate hygiene. With this increase burden of nosocomial infection it is becoming difficult for infection control committees to control the infection. However, World Health Organization gave a

standard and systematic procedure which should be followed by every hospital staff to control the nosocomial infection. Every individual of hospital is equally responsible in reducing the risk by practicing healthy ways designed by infection control committees. With this all the hospital staff must be given proper training for controlling the infection if suspected.



References:

- Bassert, Joanna M., et al. *McCurnins Clinical Textbook for Veterinary Technicians*. Elsevier, 2018.
- Ducel, G., et al. "Prevention of Hospital-Acquired Infections: A Practical Guide." *World Health Organization*, Available at: apps.who.int/medicinedocs/documents/s16355e/s16355e.pdf.
- Fang, Qing. "Systematic Review of the Effectiveness of Healthy Hospital Infrastructure to Prevent Healthcare Acquired Infections." *University of Copenhagen*, 5 Oct. 2012, Available at: www.ssi.dk/~media/Indhold/DK-dansk/Smitteberedskab/Infektionshygiejne/Diverse/Dissertation_Qing.ashx.
- Katarina, Nadova, and Rusnakova Viera. "Benefit of Active Approach to Surveillance of Hospital-Acquired Infections." *Epidemiology: Open Access*, vol. 06, no. 02, 21 Apr. 2016.
- Khan, Hassan Ahmed, et al. "Nosocomial Infections: Epidemiology, Prevention, Control and Surveillance." *Asian Pacific Journal of Tropical Biomedicine*, vol. 7, no. 5, May 2017, pp. 478–482.
- Khan, Hassan Ahmed, et al. "Nosocomial Infections: Epidemiology, Prevention, Control and Surveillance." *Asian Pacific Journal of Tropical Biomedicine*, vol. 7, no. 5, 2017, pp. 478–482.
- Mohammed, Maazuddin, et al. "Nosocomial Infections: An Overview." *International Research Journal of Pharmacy*, vol. 5, no. 1, 2014, pp. 7–12.
- Mourud, Heidi Riddervold. "Prevention of Nosocomial Infections." *Charles University in Prague*, 2010. Available at: dspace.cuni.cz/bitstream/handle/20.500.11956/30027/DPTX_2009_1_11120_0_45014_0_76768.pdf?sequence=1.
- Nazir, Asifa, and S Kadri. "An Overview of Hospital Acquired Infections and the Role of the Microbiology Laboratory." *International Journal of Research in Medical Sciences*, vol. 2, no. 1, Feb. 2014, pp. 21–27.

Nosocomial Infections: Literature Review.” *UKEssays*, 23 Mar. 2015, Available at: www.ukessays.com/essays/health-and-social-care/nosocomial-infections-review-of-literature-health-and-social-care-essay.php.

Özer, Türkan Toka, et al. “Nosocomial Infections in a District Hospital in Turkey.” *Biomedical Research: An International Journal of Medical Sciences*, vol. 26, no. 2, 2015.

Patel, Beena, et al. “Study of Microorganisms Associated with Hospital Acquired Infection in Intensive Care Unit and Their Antibiotic Resistance Pattern.” *International Journal of Scientific Research*, Vol. 2, no. 6, June 2013, pp. 447–450.

Root, Richard K., et al. *Clinical Infectious Diseases: a Practical Approach*. Oxford Univ. Press, 1999.

Sharma, Sumit. “The Menace of Nosocomial Infections.” *Global Journal of Otolaryngology*, vol. 7, no. 4, 5 May 2017.